

## Anatomy

# Visual Cortex

The visual cortex<sup>1)</sup> is a part of the brain that is responsible for processing visual information. It is located in the back of the brain, in an area called the occipital lobe.

The visual cortex is divided into several different areas, each of which is responsible for processing different types of visual information. Some of the main areas of the visual cortex include:

- **The primary visual cortex (V1):** This is the first area of the brain to receive input from the retina. It is responsible for processing basic visual features such as color, contrast, and form.
- **The secondary visual cortex (V2):** This area is responsible for processing more complex visual information, such as the relationships between objects in a scene.
- **The tertiary visual cortex (V3):** This area is involved in processing information about the spatial relationships between objects in a scene and their distance from the viewer.
- **The quaternary visual cortex (V4):** This area is involved in processing information about color and form in more complex scenes.
- **V5 is also known as the middle temporal area (MT)** and is involved in the processing of visual information related to motion. It is particularly important for detecting and tracking moving objects.
- **V6** is involved in the processing of visual information related to three-dimensional space and depth perception. It is also involved in the control of eye movements.
- **V7** is involved in the processing of visual information related to surface orientation and the perception of shape.

The visual cortex is essential for our ability to see and understand the world around us. It receives input from the retina and processes this information to create a coherent visual experience. This processed information is then passed on to other areas of the brain for further processing and integration with other senses.

## Visual Fields

The visual field is the area that can be seen by an eye (or both eyes) while the gaze is fixed in one position. It is the portion of the environment that is visible to an individual at any given moment.

The visual field can be divided into two parts: the central visual field and the peripheral visual field. The central visual field is the area of the visual field that is directly in front of the viewer and is the part of the visual field that is used for most detailed tasks, such as reading and writing. The peripheral visual field is the area of the visual field that is outside of the central visual field and is used for detecting movement and perceiving the overall layout of the environment.

The visual field is an important aspect of vision and is used to gather information about the environment. It is also important for maintaining balance and orientation in space. Visual field defects

can be caused by a variety of factors, including damage to the eye or the visual pathways in the brain, and can lead to visual impairments.

## Blindsight

Blindsight is a phenomenon that refers to the ability of some people with damage to the primary visual cortex (V1) to respond to visual stimuli, despite not being consciously aware of seeing them. Blindsight is an important phenomenon because it demonstrates the distinction between conscious perception and unconscious processing of visual information.

The primary visual cortex, also known as the striate cortex, is the first stage in the processing of visual information in the brain. It is responsible for the initial processing of visual features such as color, form, and movement. Damage to the primary visual cortex can result in a condition known as cortical blindness, where the individual is unable to consciously perceive visual stimuli in the affected part of their visual field.

However, despite the loss of conscious perception, individuals with blindsight are still able to respond to visual stimuli in an unconscious manner. For example, they may be able to accurately reach out and touch an object, or make a saccade (a rapid eye movement) towards a visual stimulus, without being consciously aware of seeing it. This suggests that there are other regions of the brain, such as the extrastriate cortex, that can process visual information even if the primary visual cortex is damaged.

The exact mechanisms underlying blindsight are still not well understood. One theory is that the extrastriate cortex is able to process visual information in an unconscious manner and that this information is then sent to the brainstem, which it can use to guide movements and reflexes. Another theory is that the unconscious processing of visual information is facilitated by the connections between the primary visual cortex and subcortical regions, such as the superior colliculus, which is involved in the control of eye movements and attention.

Regardless of the exact mechanisms, blindsight highlights the importance of considering the distinction between conscious and unconscious processing of visual information. It also has implications for our understanding of the neural basis of perception and the role of different regions of the brain in visual processing.

There are several famous cases of blindsight that have been studied and documented by neuroscientists and physicians. Some of the most well-known examples include:

- **DF:** DF is a patient who suffered damage to the primary visual cortex as a result of a stroke. Despite being unable to consciously perceive visual stimuli in one part of her visual field, she was still able to accurately reach out and touch objects, and make accurate saccades (rapid eye movements) towards stimuli.
- **TN:** TN is another patient who suffered damage to the primary visual cortex as a result of a stroke. Despite being unable to consciously perceive visual stimuli, he was still able to accurately identify the orientation of lines and the location of objects in his blind field.
- **SM:** SM is a patient who suffered damage to the primary visual cortex due to a congenital condition. Despite being unable to consciously perceive visual stimuli, he was still able to accurately reach out and touch objects, and make accurate saccades towards stimuli.

These cases and others like them have helped to shed light on the mechanisms underlying blindsight, and have contributed to our understanding of the neural basis of perception. They also have implications for our understanding of the brain and the relationship between different regions in the processing of visual information.

## Related Conditions

There are several related conditions that can arise from specific damage to the visual cortex, including:

- **Cortical blindness:** Cortical blindness is a condition that results from damage to the primary visual cortex (V1), causing a loss of conscious perception of visual stimuli. Individuals with cortical blindness may have some residual visual function, such as light perception, but are unable to consciously see anything in their affected visual field.
- **Hemianopia:** Hemianopia is a condition that results from damage to one-half of the visual cortex, causing a loss of vision in half of the visual field. Hemianopia can be caused by a variety of conditions, including stroke, brain tumors, and head injury.
- **Prosopagnosia:** Prosopagnosia, also known as face blindness, is a condition that results from damage to the fusiform gyrus, a region of the brain involved in recognizing faces. Individuals with prosopagnosia are unable to recognize faces, even though they can recognize other objects and have normal vision.
- **Balint syndrome:** Balint syndrome is a condition that results from damage to the parietal lobe, a region of the brain involved in attention and visual processing. Individuals with Balint syndrome may have difficulty with visual attention, visuospatial orientation, and visual memory.
- **Visual agnosia:** Visual agnosia is a condition that results from damage to the ventral stream, a pathway in the brain involved in object recognition. Individuals with visual agnosia are unable to recognize objects, even though they have normal vision.

These conditions demonstrate the importance of the different regions of the brain in visual processing and highlight the complex relationship between different regions in the processing of visual information. They also have implications for our understanding of the neural basis of perception and the mechanisms underlying different aspects of visual processing.

## Prosopagnosia

Prosopagnosia, also known as face blindness, is a neuropsychological disorder that affects an individual's ability to recognize faces. This condition is caused by damage to the fusiform gyrus, a region of the brain involved in recognizing faces. Prosopagnosia can be congenital or acquired and is characterized by difficulty recognizing familiar faces, including those of family members and close friends. In this essay, we will discuss the causes, symptoms, and neurological aspects of prosopagnosia.

**Causes:** Prosopagnosia can be congenital or acquired. Congenital prosopagnosia is believed to be caused by a genetic factor and is present from birth. Acquired prosopagnosia is caused by damage to the fusiform gyrus, which can result from a variety of conditions, including stroke, traumatic brain injury, and neurodegenerative diseases such as Alzheimer's disease.

**Symptoms:** Individuals with prosopagnosia experience difficulty recognizing faces, even those of close family members and friends. This difficulty can result in social difficulties, as the individual may not be able to recognize familiar faces in social situations. Individuals with prosopagnosia may also experience difficulty recognizing other facial features, such as hair and eye color, and may have trouble with facial expressions and emotions.

**Neurological aspects:** Prosopagnosia is believed to be caused by damage to the fusiform gyrus, a region of the brain involved in recognizing faces. This region is specialized for processing facial features and is highly interconnected with other regions of the brain involved in visual processing, memory, and attention. When the fusiform gyrus is damaged, the ability to recognize faces is impaired, resulting in prosopagnosia.

Studies using functional magnetic resonance imaging (fMRI) have shown that individuals with prosopagnosia have decreased activity in the fusiform gyrus when viewing faces, compared to individuals without the condition. This suggests that the fusiform gyrus plays a critical role in the recognition of faces, and that damage to this region can result in prosopagnosia.

## Fusiform Gyrus

The fusiform gyrus is a region of the brain located in the temporal lobe, and is responsible for processing and recognizing faces, as well as other complex visual stimuli. This region is thought to be one of the primary centers in the brain for face recognition, and damage to the fusiform gyrus can result in prosopagnosia, a condition characterized by difficulty recognizing faces.

The fusiform gyrus is part of the ventral visual pathway, which is involved in object recognition and perception. This pathway is important for processing information about the shape, color, and texture of objects, and for allowing us to identify and categorize different objects in our environment. The fusiform gyrus is specialized for processing facial features, such as the eyes, nose, and mouth, and is thought to have a highly interconnected network of neurons that work together to recognize faces.

Studies using functional magnetic resonance imaging (fMRI) have shown that activity in the fusiform gyrus is strongly correlated with the perception of faces. When viewing faces, the fusiform gyrus is one of the most active regions in the brain, and its activity increases with the complexity of the face being viewed. This suggests that the fusiform gyrus is critical for the recognition of faces and other complex visual stimuli.

In addition to processing facial features, the fusiform gyrus is also involved in the perception of other complex visual stimuli, such as letters and numbers, animals, and objects. This region is thought to be involved in the development of visual expertise, such as the ability to recognize faces and other objects with high accuracy.

~~socialite~~

---

<sup>1)</sup> Visual cortex [Wikipedia](#)

From:

<https://dokuwiki.3dd.de/> - **Integral Eye Movement Therapy (IEMT) Wiki**

Permanent link:

[https://dokuwiki.3dd.de/visual\\_cortex?rev=1678039139](https://dokuwiki.3dd.de/visual_cortex?rev=1678039139)

Last update: **2023/03/05 18:58**

